SNELLEN METRIC TEST CHART
Visual acuity 6/6 (20/20) Chart no. 2

SNELLEN METRIC TEST CHART
Visual acuity 6/6 (20/20) Chart no. 2

I 49.89/2: 85 (15) map 37

ORIGINAL

UNITED STATES
DEPARTMENT OF THE INTERIOR
GEOLOGICAL SURVEY

PASS-A-GRILLE BEACH QUADRANGLE
FLORIDA
7.5 MINUTE SERIES (TOPOGRAPHIC)



MAP B7

TAMPA BAY ENVIRONMENTAL ATLAS

WATER RESOURCES

PASS-A-GRILLE







Scale 1:24,000

1984

TOPICAL MAPS IN THIS ATLAS:
A. BIOLOGICAL RESOURCES
B. WATER RESOURCES

Produced by U.S. Fish and Wildlife Service
Projection and 1,000 meter grid, zone 17, Universal Transverse Mercator
10,000-foot grid ticks based on Florida coordinate system, east and west zones

Overlay material compiled in 1984 from latest available data obtained from



Produced by U.S. Fish and Wildlife Service

Projection and 1,000 meter grid, zone 17, Universal Transverse Mercator
10,000-foot grid ticks based on Florida coordinate system, east and west zones

Overlay material compiled in 1986 from latest available data obtained from Federal, State and Local government agencies and other sources.

There is no map A19 due to a lack of data.

LEGEND

35 — MEAN ANNUAL MID-DEPTH SPECIFIC CONDUCTIVITY ISOPLETH (WATER QUALITY) CONTOUR MAP
Five year mean annual specific conductivity September 1978 - August 1983 (mmhos)

POINT SOURCE DISCHARGES *

▲10 Q=___ Industrial Point Source Discharge: D is permitted capacity in total gallons per day (tgd) or millions of gallons per day (mgd).

△10 Q=___ Municipal Point Source Discharge: D is permitted capacity in total gallons per day (tgd) or millions of gallons per day (mgd).

DREDGE SPOIL DISPOSAL SITES

TIDE STATION *
(National Ocean Service)

10 Active tide station. The tide station number is listed next to the symbol.

10 Inactive tide station location. The tide station number is listed next to the symbol.

WATER QUALITY SAMPLING STATION *


TAMPA BAY
NATIONAL WILDLIFE REFUGE
Mullet Key
Mullet Key Bayou
Fort Desoto County Park
Egmont Key
Egmont Key Bayou
Shell Key
Middle Ground
Mullet Key Shoal
Mullet Key Channel
Tampa Bay Entrance Channel
Anchorage
Anchorage Area
Fish Haulout
Manatee Haulout
Bridge Span Bank
Dumfoundling Bay
Dumfoundling Key
Eastern Spit
Mule Key
Whale Island
Hurricane Ridge
Middle Ground Shoal
Tampa Bay Entrance Channel
Mullet Key Channel
Mullet Key Bayou
Egmont Key Bayou
Egmont Key
Shell Key
Fort Desoto County Park
Fish Haulout
Manatee Haulout
Bridge Span Bank
Anchorage
Anchorage Area

THE TAMPA BAY ENVIRONMENTAL ATLAS WAS PRODUCED BY THE U.S. FISH AND WILDLIFE SERVICE, DIVISION OF BIOLOGICAL SERVICES NATIONAL COASTAL ECOSYSTEMS TEAM (FWS/OBS) FOR COASTAL PLANNERS, DECISION MAKERS, AND THE GENERAL PUBLIC. TO DETERMINE AND EVALUATE CHANGE AND TO ASSESS THE EFFECTS OF MAN'S ACTIVITIES ON THE QUALITY OF ESTUARIES AND THE MAINTENANCE OF ECOLOGICAL DIVERSITY. THE STUDY AREA INCLUDES A MAJOR PORTION OF THE COASTAL COUNTIES OF PINELLAS, HILLSBOROUGH, MANATEE, AND SARASOTA, FLORIDA. THE ATLAS IS NOT A PLANNING DOCUMENT OR PRODUCT OF THIS STUDY. TO DEFINE THE LIMITS OF PROPRIETARY JURISDICTION OF ANY FEDERAL,STATE, OR LOCAL GOVERNMENT TO ESTABLISH THE GEOGRAPHICAL SCOPE OF THE REGULATORY PROGRAMS OF GOVERNMENT AGENCIES.

THIS MAP HAS BEEN COMPILED FROM A VARIETY OF SOURCE MATERIALS, INCLUDING MAPS, AERIAL PHOTOS, AND REPORTS. SINCE MANY SOURCES CONTAINED GENERALIZED LOCATIONS OF RESOURCES AND OTHER MAPPED FEATURES, THE USER SHOULD NOT RELY ON THESE MAPS FOR PRECISE MEASUREMENTS OF DISTANCES OR AREAS.

PRIMARY SOURCES OF MAPPED INFORMATION

SALINITY: Conductivity for Tampa Bay water quality sampling program, 1972-1980. Tampa: Hillsborough County Environmental Protection Commission.

POINT SOURCE DISCHARGES: Composer grimmicyl of industrial- and municipal point source discharges in West Central Florida. Florida Dept. of Environmental Regulation,1983.

DREDGE SPOIL DISPOSAL AREAS: Dredge and Spoil easement location maps. Florida Dept. of Natural Resources

N.O.S. TIDE STATIONS: Index of tide stations for the state of Florida. National Ocean Service, Rockville, Md. 1981

WATER QUALITY: Total chlorophyll, conductivity, turbidity, and temperature data collected at stations in the Tampa Bay water quality sampling. Sept. 1979 - August 1983. Hillsborough County Environmental Protection Commission.

BATHYMETRIC: Bathymetric map by St. Petersburg Quadrangle, N.O.S. 1978.

INTERTIDAL ZONE: Nautical Chart #'s 11410, 11412, 11414 and 11420, N.O.S., Rockville, Md. 1982.

SEDIMENTS: Sediments, oceanographic observations and data. Tampa Bay, Florida. Vol. 1 and 2. Bottom sediment sation, 1981-1985. Taylor and Salomon, 1985

TIDAL CURRENTS: Changes in tidal flow, circulation and flushing caused by dredge and fill in Tampa Bay, Florida U.S. Geological Survey Open File Report 85-847 Goodwin, 1985

FRESHWATER-SALTWATER INTERFACE: Position of the Freshwater- Saltwater interface in the upper part of the Floridan aquifer, Southwest Florida, 1979. U.S.G.S. water resources investi- tion report 82-30. Causseaux and Fretwell, 1982

FOR COMPLETE REFERENCE SEE MAP NARRATIVE

SCALE 1:24000

0 1000 0 2000 3000 4000 5000 6000 7000 FEET

0.5 0 1 0 2 KILOMETER



Note: Specific conductivity values on the supplemental legend should be in pmhos.

Supplemental Legend for Water Quality Stations on This Map

Station #25 — Lat. 27°28'13" — Long. 82°40'13"

PARAMETER	SPRING (3-5)	SUMMER (6-8)	FALL (9-11)	WINTER (12-2)	ANNUAL (1-12)
Total Chlorophyll (ug/l)	4.0	4.0	5.1	3.0	4.0
Conductivity (mmhos)	4983	4765	4769	5186	4926
Turbidity (FTU'S)	4.8	4.0	5.1	3.0	4.4
Temperature (°C)	23.9	30.0	25.0	18.8	23.9
Total Nitrogen (mg/l)	0.51	0.53	0.65	0.39	0.52

Station #50 — Lat. 27°28'19" — Long. 82°41'18"

PARAMETER	SPRING (3-5)	SUMMER (6-8)	FALL (9-11)	WINTER (12-2)	ANNUAL (1-12)
Total Chlorophyll (ug/l)	17.0	19.7	22.2	14.4	18.5
Conductivity (mmhos)	51167	49944	49851	52278	49310
Turbidity (FTU'S)	5.8	3.8	7.1	4.8	5.2
Temperature (°C)	22.8	30.1	25.2	18.4	23.6
Total Nitrogen (mg/l)	0.45	0.85	0.85	0.51	0.62

Current Velocity

Resultant tidal current. The direction is indicated by the arrow and the discharge range is feet per second

- 0.01 - 0.10 ft/sec
- 0.11 - 0.20 ft/sec
- 0.21 - 0.30 ft/sec
- 0.31 - 0.50 ft/sec
- >=0.50 ft/sec

FRESHWATER/SALTWATER INTERFACE

(U.S. Geological Survey 1979)
250 mg/l line of equal chloride concentration in the upper part of Floridan aquifer.

**Symbols appearing sum numbers are described more fully in the narrative*

MICROFILMED FROM
BEST AVAILABLE COPY

MICROCOPY RESOLUTION TEST CHART
NATIONAL BUREAU OF STANDARDS-1963-A
MICROFILMED FROM
BEST AVAILABLE COPY